

# Recombinant Mouse Fibroblast Growth Factor-acidic

## Mouse, Recombinant (Fgf1)

Expressed in *E. coli*

Cat. No. CRP0828

Lot. No. (See product label)

### PRODUCT INFORMATION

**Description:** FGF acidic, also known as FGF-1, ECGF, and HBGF-1, is a 17 kDa nonglycosylated member of the FGF family of mitogenic peptides. FGF acidic, which is produced by multiple cell types, stimulates the proliferation of all cells of mesodermal origin and many cells of neuroectodermal, ectodermal, and endodermal origin. It plays a number of roles in development, regeneration, and angiogenesis. FGF-acidic is a non-glycosylated heparin binding growth factor that is expressed in the brain, kidney, retina, smooth muscle cells, bone matrix, osteoblasts, astrocytes and endothelial cells. FGF-acidic has the ability to signal through all the FGF receptors.

**Amino-Acid Sequence:** 140aa (The sequence of the first five N-terminal amino acids was determined and was found to be Phe-Asn-Leu-Pro-Leu.), non-glycosylated.

**M. W. :** 15,796 Da

**Recombinant:** Expressed in *E. coli*

**Purity:** >95% as determined by HPLC and SDS-PAGE .


**Formulation:** Mouse FGF-acidic was lyophilized after extensive dialysis against PBS.

**Specific Activity:** The ED50 as determined by the dose-dependant stimulation of thymidine uptake by 3T3 cells in the presence of heparin was found to be less than 0.5 ng/ml, corresponding to a Specific Activity of  $2.0 \times 10^6$  IU/mg.

**Endotoxin:** Less than 0.1ng/μg (1 IEU/μg) of FGF-acidic.

**Reconstitution:** It is recommended to reconstitute the lyophilized rmFGF-acidic in sterile 18MΩ-cm H<sub>2</sub>O not less than 100μg/ml, which can then be further diluted to other aqueous solutions.

**Storage:** Lyophilized rmFGF-acidic although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution rmFGF-acidic should be stored at 4°C between 2-7 days and for future use below -18°C. For long-term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Aliquot to avoid repeated freeze-thaw cycles.



[PDB](#) rendering based on 1afc.

### GENE INFORMATION

**Gene Name:** [FGF1](#)

**Synonyms:** Dffrx, Fam, Fgf-1, Fgfa, HBGF-1, ECGF-beta, FIBP, FGFIBP, FIBP-1, ECGF, ECGFA, GLIO703, FGF1, FGF-a, Heparin-binding growth factor 1, Acidic fibroblast growth factor, aFGF; ECGFB; FGF-alpha; FGFA; HBGF1; AFGF; Beta-endothelial cell growth factor; Heparin binding growth factor 1 precursor; endothelial cell growth factor alpha; endothelial cell growth factor beta; fibroblast growth factor 1 (acidic); heparin binding growth factor 1

**mRNA Refseq:** [NM\\_010197](#)

**Protein Refseq:** [NP\\_034327](#)

**MIM:** [131220](#)

**GeneID:** [14164](#)

**UniProt ID:** P61148

**Chromosome Location:** 18 19.0 cM

**Pathway:** MAPK signaling pathway; Melanoma; Regulation of actin cytoskeleton

**Function:** growth factor activity; heparin binding

### REFERENCES

1.Yu YL, Kha H, Golden JA, et al. An acidic fibroblast growth factor protein generated by alternate splicing acts like an antagonist. *J. Exp. Med.* 1992; 175 (4): 1073–1080

**FOR RESEARCH USE ONLY**

Creative Biolabs All rights reserved.

45-16 Ramsey Road Shirley, NY 11967, USA  
Technical Support: T: 631-871-5806 · F: 631-207-8356  
E-mail: [info@creative-biolabs.com](mailto:info@creative-biolabs.com)  
[www.creative-biolabs.com](http://www.creative-biolabs.com)