

Recombinant Human Interferon-alpha 2b

Human, Recombinant (INFA2)

Expressed in *E. coli*

Cat. No. CRP0813

Lot. No. (See product label)

PRODUCT INFORMATION

Description: Recombinant Human Interferon-alpha 2b produced in *E. coli* is a single, non-glycosylated, polypeptide chain containing 166 amino acids and having a molecular mass of 19400 Dalton. The Interferon-alpha 2b gene was obtained from human leukocytes.

Amino-Acid Sequence: 166 aa (The sequence of the first fifteen N-terminal amino acids was determined and was found to be Met-Cys-Asp-Leu-Pro-Gln-Thr-His-Ser-Leu-Gly-Ser-Arg-Arg-Thr-Leu.), non-glycosylated

M. W. : 19,200 Da

Recombinant: Expressed in *E. coli*

Purity: >95% as determined by SEC-HPLC and SDS-PAGE .

Formulation: Lyophilized from a (1mg/ml) solution in containing 8.8mg sodium chloride, 0.4mg sodium acetate and 1.0mg acetate acid buffer.

Specific Activity: The specific activity as determined in a viral resistance assay using VSV- WISH cells was found to be greater than 1.0×10^8 IU/ mg.

Endotoxin: Less than 0.3ng/μg (0.3IEU/μg) determined by LAL test.

Reconstitution: It is recommended to reconstitute the lyophilized rHuIFN-alpha2b in sterile 18MΩ-cm H₂O not less than 100μg/ml, which can then be further diluted to other aqueous solutions.

Storage: Lyophilized rHuIFN-alpha 2b although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution rHuIFN-alpha 2b should be stored at 4°C between 2-7 days and for future use below -18°C. For long-term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Aliquot to avoid repeated freeze-thaw cycles.

FOR RESEARCH USE ONLY

GENE INFORMATION

Gene Name: [IFNA2](#)

Synonyms: IFNA, MGC125764, MGC125765, Leukocyte interferon, B cell interferon, Type I interferon, IFN-α 2a, Interferon alpha-A, LelF A, Interferon alpha-2, Interferon alpha-2 Precursor

mRNA Refseq: [NM_000605.2](#)

Protein Refseq: [NP_000596.2](#)

MIM: [147562](#)

GeneID: [3440](#)

UniProt ID: P01563

Chromosome Location: 9p22

Pathway: Antigen processing and presentation; Cytokine-cytokine receptor interaction ;Jak-STAT signaling pathway ;KEGG pathway: Natural killer cell mediated cytotoxicity ;Regulation of autophagy ;Toll-like receptor signaling pathway

Function: interferon-alpha/beta receptor binding

REFERENCES

- 1.Thomas H. et al. Mechanisms of Action of Interferon and Nucleoside Analogues. Journal of hepatology.2004 Feb; 40 (2): 364
- 2.Schmeisser H, Gorshkova I, et al. Two interferons alpha influence each other during their interaction with the extracellular domain of human type interferon receptor subunit 2. Biochemistry. 2007 Dec 18; 46(50):14638-14649.
- 3.Thyrell et al. Mechanisms of Interferon-alpha induced apoptosis in malignant cells. Oncogene: Nature Publishing Group. 2002; 21:1251-1262

Creative Biolabs All rights reserved.

45-16 Ramsey Road Shirley, NY 11967, USA
Technical Support: T: 631-871-5806 ·F: 631-207-8356
E-mail: info@creative-biolabs.com
www.creative-biolabs.com