

# Recombinant Human Tumor Necrosis Factor alpha-1a

Human, Recombinant (TNF)

Expressed in *E. coli*

Cat. No. CRP0806

Lot. No. (See product label)

## PRODUCT INFORMATION

**Description:** TNF-alpha is a homotrimer with a subunit molecular mass of 17 kDa and that it plays a major role in growth regulation, differentiation, inflammation, viral replication, tumorigenesis, and autoimmune diseases; and in viral, bacterial, fungal, and parasitic infections. Besides inducing hemorrhagic necrosis of tumors, TNF was found to be involved in tumorigenesis, tumor metastasis, viral replication, septic shock, fever, inflammation, and autoimmune diseases including Crohn's disease, and rheumatoid arthritis as well as graft-versus-host disease.

**Amino-Acid Sequence:** 158 aa(The sequence of the first fifteen N-terminal amino acids was determined and was found to be Met-Val-Arg-Ser-Ser-Ser-Arg-Thr-Pro-Ser-Asp-Lys-Pro-Val-Ala), non-glycosylated

**M. W. :** 17,484 Da

**Recombinant:** Expressed in *E. coli*

**Purity:** >95% as determined by SDS-PAGE and SEC-HPLC.

**Isoelectric Point:** The main zone between 4.0~5.0 analysis by IEF.

**UV scan:** The maximal absorption wave is 275+/-3nm.


**Formulation:** Lyophilized from a 0.2 µm filtered solution in PBS.

**Specific Activity:** The ED50 as determined by the cytolysis of murine L929 cells in the presence of Actinomycin D is less than 0.03ng/ml, corresponding to a Specific Activity of  $3.0 \times 10^7$  IU/mg.

**Endotoxin:** Less than 0.01ng/µg (0.01IEU/µg) determined by LAL test.

**Reconstitution:** It is recommended to reconstitute the lyophilized TNF alpha-1a in sterile 18MΩ-cm H<sub>2</sub>O not less than 100µg/ml, which can then be further diluted to other aqueous solutions.

**Storage:** Lyophilized samples are stable for up to twelve months from date of receipt at -20°C to -70°C.



[PDB](#) rendering based on 1TNF.

## GENE INFORMATION

**Gene Name:** [TNF](#)

**Synonyms:** tumor necrosis factor (TNF superfamily, member 2); DIF; TNFA; TNFSF2; TNF-alpha; Cachectin; OT-THUMP00000037669; TNF-a; APC1 protein; TNF superfamily, member 2; TNF, macrophage-derived; TNF, monocyte-derived; Tumor necrosis factor ligand superfamily member 2; Tumor necrosis factor precursor; tumor necrosis factor; tumor necrosis factor (cachectin); tumor necrosis factor alpha; tumor necrosis factor, alpha (cachectin); TNF

**mRNA Refseq:** [NM\\_000594](#)

**Protein Refseq:** [NP\\_000585](#)

**MIM:** [191160](#)

**UniProt ID:** P01375

**Gene ID:** [7124](#)

**Chromosome Location:** 6p21.3

**Function:** cytokine activity, identical protein binding, tumor necrosis factor receptor binding

## REFERENCES

- 1.Gaur U, Aggarwal BB (2003).Regulation of proliferation, survival and apoptosis by members of the TNF superfamily. *Biochem. Pharmacol.* 66 (8): 1403–1408
- 2.Wajant H, Pfizenmaier K, Scheurich P (2003). Tumor necrosis factor signaling. *Cell Death Differ.* 10 (1): 45–65
- 3.Tang P, Hung M-C, Klostergaard J (1996). Human pro-tumor necrosis factor is a homotrimer. *Biochemistry* 35 (25): 8216–8225

**FOR RESEARCH USE ONLY**

Creative Biolabs All rights reserved.

45-16 Ramsey Road Shirley, NY 11967, USA  
Technical Support: T: 631-871-5806 ·F: 631-207-8356  
E-mail: [info@creative-biolabs.com](mailto:info@creative-biolabs.com)  
[www.creative-biolabs.com](http://www.creative-biolabs.com)