

Product Information

Recombinant Anti-Human ERBB2 Antibody scFv Fragment

Cat. No.: **MOM-18005-S(P)**

This product is for research use only and is not intended for diagnostic use.

Product Overview

Recombinant Humanized (from mouse) Antibody scFv Fragment is specific to Human ErbB2, expressed in E. coli

Antigen Description

Upon ERBB2 activation, the MEMO1-RHOA-DIAPH1 signaling pathway elicits the phosphorylation and thus the inhibition of GSK3B at cell membrane. This prevents the phosphorylation of APC and CLASP2, allowing its association with the cell membrane. In turn, membrane-bound APC allows the localization of MACF1 to the cell membrane, which is required for microtubule capture and stabilization. In the nucleus is involved in transcriptional regulation. Associates with the 5'-TCAAATTC-3' sequence in the PTGS2/COX-2 promoter and activates its transcription. Implicated in transcriptional activation of CDKN1A; the function involves STAT3 and SRC.

Specific Activity

Tested positive against native antigen.

Target

ErbB2

Immunogen

NIH 3T3/HER2-3400 cells are used to immunize BALB/c mice.

Source

Humanized (from mouse)

Species Reactivity

Human

Type

scFv Fragment from Humanized (from mouse) IgG1 - kappa

Expression Host

E. coli

Purity

>95.0% as determined by Analysis by RP-HPLC & analysis by SDS-PAGE.

Applications

Suitable for use in ELISA, WB, Neut and most other immunological methods.

Storage

Store at -20°C for long-term storage. Store at 2-8°C for up to one month. Avoid freeze/thaw cycles.

ANTIGEN GENE INFORMATION

Gene Name

[ERBB2 v-erb-b2 erythroblastic leukemia viral oncogene homolog 2, neuro/glioblastoma derived oncogene homolog \(avian\) \[Homo sapiens \]](#)

Official Symbol

ERBB2

Synonyms

ERBB2; v-erb-b2 erythroblastic leukemia viral oncogene homolog 2, neuro/glioblastoma derived oncogene homolog (avian); NGL, v erb b2 avian erythroblastic leukemia viral oncogene homolog 2 (neuro/glioblastoma derived oncogene homolog); receptor tyrosine-protein kinase erbB-2; CD340; HER 2; HER2; NEU; herstatin; p185erbB2; proto-oncogene Neu; c-erb B2/neu protein; proto-oncogene c-ErbB-2; metastatic lymph node gene 19 protein; tyrosine kinase-type cell surface receptor HER2; neuroblastoma/glioblastoma derived oncogene homolog; v-erb-b2 avian erythroblastic leukemia viral oncogene homolog 2 (neuro/glioblastoma derived oncogene homolog); NGL; TKR1; HER-2; MLN 19; HER-2/neu;

Gene ID

[2064](#)

mRNA Refseq

[NM_001005862](#)

Protein Refseq

[NP_001005862](#)

MIM

[164870](#)

UniProt ID

P04626

Chromosome Location

17q11.2-q12

Pathway

Adherens junction, organism-specific biosystem; Adherens junction, conserved biosystem; Alpha6-Beta4 Integrin Signaling Pathway, organism-specific biosystem; Axon guidance, organism-specific biosystem; Bladder cancer, organism-specific biosystem; Bladder cancer, conserved biosystem; Calcium signaling pathway, organism-specific biosystem;

Function

ATP binding; ErbB-3 class receptor binding; Hsp90 protein binding; RNA polymerase I core binding; epidermal growth factor-activated receptor activity; glycoprotein binding; contributes_to growth factor binding; identical protein binding; nucleotide binding; protein C-terminus binding; protein binding; protein dimerization activity; protein heterodimerization activity; protein heterodimerization activity; protein heterodimerization activity; protein phosphatase binding; protein tyrosine kinase activity; protein tyrosine kinase activity; protein tyrosine kinase activity; receptor activity; receptor signaling protein tyrosine kinase activity; transmembrane receptor protein tyrosine kinase activity; transmembrane signaling receptor activity; ubiquitin protein ligase binding;