

Product Information

Recombinant Anti-Human MUC1 Antibody Fab Fragment

Cat. No.: **MOM-18037-F(P)**

This product is for research use only and is not intended for diagnostic use.

Product Overview

Recombinant Humanized (from mouse) Antibody Fab Fragment is bind to Human MUC1, expressed in E. coli

Antigen Description

The alpha subunit has cell adhesive properties. Can act both as an adhesion and an anti-adhesion protein. May provide a protective layer on epithelial cells against bacterial and enzyme attack. The beta subunit contains a C-terminal domain which is involved in cell signaling, through phosphorylations and protein-protein interactions. Modulates signaling in ERK, SRC and NF-kappa-B pathways. In activated T-cells, influences directly or indirectly the Ras/MAPK pathway. Promotes tumor progression. Regulates TP53-mediated transcription and determines cell fate in the genotoxic stress response. Binds, together with KLF4, the PE21 promoter element of TP53 and represses TP53 activity.

Specific Activity

Tested positive against native antigen.

Target

MUC1

Immunogen

The details of the immunogen for this antibody are not available.

Source

Humanized (from mouse)

Species Reactivity

Human

Type

Fab Fragment based on Humanized (from mouse) Fab - G1 - kappa

Expression Host

E. coli

Predicted N terminal

H chain: QVQLQQS; L Chain: DIQLTQS

Purity

>95.0% as determined by analysis by SDS-PAGE.

Applications

Suitable for use in FC, IP, ELISA, Neut, FuncS, IF and most other immunological methods.

Storage

Store at -20°C for long-term storage. Store at 2-8°C for up to one month. Avoid freeze/thaw cycles.

ANTIGEN GENE INFORMATION

Gene Name

[MUC1 mucin 1, cell surface associated \[Homo sapiens \]](#)

Official Symbol

MUC1

Synonyms

MUC1; mucin 1, cell surface associated; mucin 1, transmembrane , PUM; mucin-1; CD227; PEM; episialin; DF3 antigen; H23 antigen; krebs von den Lungen-6; mucin 1, transmembrane; tumor-associated mucin; carcinoma-associated mucin; polymorphic epithelial mucin; peanut-reactive urinary mucin; tumor associated epithelial mucin; breast carcinoma-associated antigen DF3; tumor-associated epithelial membrane antigen; EMA; PUM; KL-6; MAM6; PEMT; H23AG; MUC-1; MUC-1/X; MUC1/ZD; MUC-1/SEC;

Gene ID

[4582](#)

mRNA Refseq

[NM_001018016](#)

Protein Refseq

[NP_001018016](#)

MIM

[158340](#)

UniProt ID

P15941

Chromosome Location

1q22

Pathway

IL-7 Signaling Pathway, organism-specific biosystem; Metabolism of proteins, organism-specific biosystem; O-linked glycosylation of mucins, organism-specific biosystem; Post-translational protein modification, organism-specific biosystem; T Cell Receptor Signaling Pathway, organism-specific biosystem; Termination of O-glycan biosynthesis, organism-specific biosystem;

Function

RNA polymerase II core promoter proximal region sequence-specific DNA binding; p53 binding; protein binding; transcription cofactor activity;