

Product Information

Recombinant Anti-Human il8 Antibody

Cat. No.: **MOM-18581**

This product is for research use only and is not intended for diagnostic use.

Product Overview

Recombinant Mouse Antibody is against Human IL8, expressed in Chinese Hamster Ovary cells(CHO)

Antigen Description

IL-8 is a chemotactic factor that attracts neutrophils, basophils, and T-cells, but not monocytes. It is also involved in neutrophil activation. It is released from several cell types in response to an inflammatory stimulus. IL-8(6-77) has a 5-10-fold higher activity on neutrophil activation, IL-8(5-77) has increased activity on neutrophil activation and IL-8(7-77) has a higher affinity to receptors CXCR1 and CXCR2 as compared to IL-8(1-77), respectively.

Specific Activity

Tested positive against native antigen.

Target

IL8

Immunogen

Recombinant full length protein (human).

Source

Mouse

Species Reactivity

Human

Type

IgG

Expression Host

CHO

Purity

Purity >95% by SDS-PAGE.

Applications

Suitable for use in Neut, ELISA, FC, IP, IHC and most other immunological methods.

Storage

Store at -20°C. Avoid multiple freeze/thaw cycles.

ANTIGEN GENE INFORMATION

Gene Name

Official Symbol

IL8

Synonyms

IL8; interleukin 8; interleukin-8; 3 10C; alveolar macrophage chemotactic factor I; AMCF I; b ENAP; beta endothelial cell derived neutrophil activating peptide; chemokine (C X C motif) ligand 8; CXCL8; GCP 1; GCP1; granulocyte chemotactic protein 1; IL 8; K60; LECT; LUCT; lung giant cell carcinoma derived chemotactic protein; lymphocyte derived neutrophil activating peptide; LYNAP; MDNCF; MONAP; monocyte derived neutrophil chemotactic factor; monocyte derived neutrophil activating peptide; NAF; NAP 1; NAP1; neutrophil activating peptide 1; SCYB8; TSG 1; tumor necrosis factor induced gene 1; emoctakin; T-cell chemotactic factor; neutrophil-activating peptide 1; chemokine (C-X-C motif) ligand 8; beta-thromboglobulin-like protein; tumor necrosis factor-induced gene 1; monocyte-derived neutrophil chemotactic factor; monocyte-derived neutrophil-activating peptide; small inducible cytokine subfamily B, member 8; lung giant cell carcinoma-derived chemotactic protein; beta endothelial cell-derived neutrophil activating peptide; GCP-1; NAP-1;

Gene ID

[3576](#)

mRNA Refseq

[NM_000584](#)

Protein Refseq

[NP_000575](#)

MIM

[146930](#)

UniProt ID

P10145

Chromosome Location

4q13-q21

Pathway

ATF-2 transcription factor network, organism-specific biosystem; Activation of Genes by ATF4, organism-specific biosystem; Amoebiasis, organism-specific biosystem; Amoebiasis, conserved biosystem; Bladder cancer, organism-specific biosystem; Bladder cancer, conserved biosystem; Calcineurin-regulated NFAT-dependent transcription in lymphocytes, organism-specific biosystem;

Function

chemokine activity; interleukin-8 receptor binding; protein binding;