

# Product Information

## **MemDX™ Antibody Discovery - Mouse ErbB3 / Her3 (20-641) Membrane Protein, Partial, -**

### **His tag**

Cat. No.: **MP0051F**

This product is for research use only and is not intended for diagnostic use.

This membrane protein is Mouse ErbB3 / Her3 (20-641). It has been tested in SDS-PAGE, ELISA. We provide this protein to facilitate your membrane protein antibody discovery and development.

### **Product Specifications**

#### **Host Species**

Mouse

#### **Target Protein**

ErbB3 / Her3

#### **Protein Length**

ECD

#### **Molecular Weight**

The protein has a calculated MW of 70.4 kDa. The protein migrates as 80-90 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

#### **Sequence**

AA Ser 20 - His 641 (Accession # Q61526-1).

### **Product Description**

#### **Activity**

Yes

#### **Application**

SDS-PAGE, ELISA

#### **Expression Systems**

HEK293

#### **Tag**

His tag at the C-terminus

#### **Protein Format**

Soluble

#### **Form**

LYOPH

### Reconstitution

Please see Certificate of Analysis for specific instructions.

### Endotoxin

<1.0 EU/μg by the LAL method

### Purity

>95% as determined by SDS-PAGE.

### Buffer

Lyophilized from 0.22 μm filtered solution in PBS, pH7.4. Normally trehalose is added as protectant before lyophilization.

### Storage

Stored at lyophilized form at -20°C or lower. Avoid repeated freeze-thaw cycles.

The antigen can be stable for 12 months in lyophilized form after storage at -20°C to -80°C, 3 months under sterile conditions after reconstitution after storage at -80°C.

## Target

### Target Protein

ErbB3 / Her3

### Full Name

erb-b2 receptor tyrosine kinase 3

### Introduction

This gene encodes a member of the epidermal growth factor receptor (EGFR) family of receptor tyrosine kinases. This membrane-bound protein has a neuregulin binding domain but not an active kinase domain. It therefore can bind this ligand but not convey the signal into the cell through protein phosphorylation. However, it does form heterodimers with other EGF receptor family members which do have kinase activity. Heterodimerization leads to the activation of pathways which lead to cell proliferation or differentiation. Amplification of this gene and/or overexpression of its protein have been reported in numerous cancers, including prostate, bladder, and breast tumors. Alternate transcriptional splice variants encoding different isoforms have been characterized. One isoform lacks the intermembrane region and is secreted outside the cell. This form acts to modulate the activity of the membrane-bound form. Additional splice variants have also been reported, but they have not been thoroughly characterized

### Alternative Names

HER; Erbb; Her3; Erbb-; C76256; Erbb-3; Erbb3r; receptor tyrosine-protein kinase erbB-3;avian erythroblastosis oncogene B 3 receptor; c-erbB-3; glial growth factor receptor; proto-oncogene-like protein c-ErbB-3; v-erb-b2 erythroblastic leukemia viral oncogene homolog 3

### Gene ID

[13867](#)

### UniProt ID

[Q61526](#)