SUITE 203, 17 Ramsey Road, Shirley, NY 11967, USA
Tel: 1-631-416-1478 Fax: 1-631-207-8356
Email: info@creative-biolabs.com
www.creative-biolabs.com

Product Information

MemDX™ Membrane Protein Human LILRB2 (Leukocyte immunoglobulin like receptor B2)

Cat. No.: **MP0178J**
This product is for research use only and is not intended for diagnostic use.

This product is a 48.6 kDa Human LILRB2 membrane protein expressed in HEK293T. The protein is for research use only and is not approved for use in humans or in clinical diagnosis.

Product Specifications

Host Species
Human

Target Protein
LILRB2

Protein Length
Full-length

Protein Class
Transmembrane

Molecular Weight
48.6 kDa

TMD
1

Sequence
MTPIVTVLICLGLSLGPRTHVQTGTIPKPTLWAEPDSVITQGSPVTLSCQGSLEAQEYRLYREKKSASWI
TRIRPELVKNGQFHIPSITWEHTGRYGCQYYSRARWSELSDPLVLVMTGAYPKPTLSAQPSPVVTSGGRV
TLQCESQVAFGGFILCKEGEDEHPQCLNSQPHARGSSRAIFSVGPVSPNRRWSHRCYGYDLNSPYVWSSP
SDLLELLVPGVSKKPSLSVQPGPVVAPGESLTLQCVSDVGYDRFVLYKEGERDLRQLPGRQPQAGLSQAN
FTLGPVSRSYGGQYRCYGAYNLSSEWSAPSDPLDILITGQIHGTPFISVQPGPTVASGENVTLLCQSWRQ
FHTFLLTKAGAADAPLRLRSIHEYPKYQAEFPMSPVTSAHAGTYRCYGSLNSDPYLLSHPSEPLELVVSG
PSMGSSPPPTGPISTPAGPEDQPLTPTGSDPQSGLGRHLGVVIGILVAVVLLLLLLLLFLILRHRRQGK
HWTSTQRKADFQHPAGAVGPEPTDRGLQWRSSPAADAQEENLYAAVKDTQPEDGVEMDTRAAASEAPQDV
TYAQLHSLTLRRKATEPPPSQEGEPPAEPSIYATLAIH

Product Description

Expression Systems
HEK293T

Tag
Tag Free

Form

Liquid

Purification

Anti-DDK affinity column followed by conventional chromatography steps

Purity

> 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer

25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10% glycerol

Storage

Store at +4°C for up to one week or several months at -80°C

Target

Target Protein

LILRB2

Full Name

Leukocyte immunoglobulin like receptor B2

Introduction

This gene is a member of the leukocyte immunoglobulin-like receptor (LIR) family, which is found in a gene cluster at chromosomal region 19q13.4. The encoded protein belongs to the subfamily B class of LIR receptors which contain two or four extracellular immunoglobulin domains, a transmembrane domain, and two to four cytoplasmic immunoreceptor tyrosine-based inhibitory motifs (ITIMs). The receptor is expressed on immune cells where it binds to MHC class I molecules on antigen-presenting cells and transduces a negative signal that inhibits stimulation of an immune response. It is thought to control inflammatory responses and cytotoxicity to help focus the immune response and limit autoreactivity. Multiple transcript variants encoding different isoforms have been found for this gene.

Alternative Names

ILT4; LIR2; CD85D; ILT-4; LIR-2; MIR10; MIR-10

Gene ID

10288

UniProt ID

A2IXV5

SUITE 203, 17 Ramsey Road, Shirley, NY 11967, USA
Tel: 1-631-416-1478 Fax: 1-631-207-8356
Email: info@creative-biolabs.com