

Product Information

MemDX™ Antibody Discovery - Human HGF R / c-MET (25-932) Membrane Protein, Partial, - hIgG1 Fc tag

Cat. No.: **MP0222F**

This product is for research use only and is not intended for diagnostic use.

This membrane protein is Human HGF R / c-MET (25-932). It has been tested in SDS-PAGE. We provide this protein to facilitate your membrane protein antibody discovery and development.

Product Specifications

Host Species

Human

Target Protein

HGF R / c-MET

Protein Length

ECD

Molecular Weight

The mature form of HGF R is a disulfide-linked heterodimer composed of proteolytically cleaved α and β chain. Each α and β chain has a calculated MW of 32.5 kDa (α chain) and 95.3 kDa (β chain Fc chimera). The protein migrates as 40-45 kDa (α chain) and 106-116 kDa (β chain Fc chimera) under reducing (R) condition (SDS-PAGE) due to glycosylation.

Sequence

AA Glu 25 - Thr 932 (Accession # AAI30421.1).

Product Description

Application

SDS-PAGE

Expression Systems

HEK293

Tag

Human IgG1 Fc tag at the C-terminus

Protein Format

Soluble

Form

LYOPH

Reconstitution

Please see Certificate of Analysis for specific instructions.

Endotoxin

<1.0 EU/µg by the LAL method

Purity

>95% as determined by SDS-PAGE.

Buffer

Lyophilized from 0.22 µm filtered solution in 50 mM Tris, 100 mM Glycine, pH7.5. Normally trehalose is added as protectant before lyophilization. Contact us for customized product form or formulation.

Storage

Stored at lyophilized form at -20°C or lower. Avoid repeated freeze-thaw cycles.

The antigen can be stable for 12 months in lyophilized form after storage at -20°C to -80°C, 3 months under sterile conditions after reconstitution after storage at -80°C.

Target

Target Protein

HGF R / c-MET

Full Name

hepatocyte growth factor

Introduction

This gene encodes a protein that binds to the hepatocyte growth factor receptor to regulate cell growth, cell motility and morphogenesis in numerous cell and tissue types. Alternative splicing results in multiple transcript variants, at least one of which encodes a preproprotein that is proteolytically processed to generate alpha and beta chains, which form the mature heterodimer. This protein is secreted by mesenchymal cells and acts as a multi-functional cytokine on cells of mainly epithelial origin. This protein also plays a role in angiogenesis, tumorogenesis, and tissue regeneration. Although the encoded protein is a member of the peptidase S1 family of serine proteases, it lacks peptidase activity. Mutations in this gene are associated with nonsyndromic hearing loss.

Alternative Names

SF; HGFB; HPTA; F-TCF; DFNB39; hepatocyte growth factor; fibroblast-derived tumor cytotoxic factor; hepatocyte growth factor (hepatopoietin A; scatter factor); hepatopoietin-A; lung fibroblast-derived mitogen

Gene ID

[3082](#)

UniProt ID

[P08581](#)