

# **Product Information**

# MemDX™ Antibody Discovery - Human Mucin-1 / MUC-1 (24-380) (24-380) Membrane Protein, Partial, -hlgG1 Fc tag

Cat. No.: MP0635F

This product is for research use only and is not intended for diagnostic use.

This membrane protein is Human Mucin-1 / MUC-1 (24-380) (24-380). It has been tested in SDS-PAGE. We provide this protein to facilitate your membrane protein antibody discovery and development.

# **Product Specifications**

# **Host Species**

Human

#### **Target Protein**

Mucin-1 / MUC-1 (24-380)

#### **Protein Length**

**ECD** 

# **Molecular Weight**

The protein has a calculated MW of 60.2 kDa.

#### Sequence

AA Ser 24 - Ser 380 (Accession # P15941-1).

#### **Product Description**

#### **Application**

SDS-PAGE

# **Expression Systems**

**HEK293** 

### Tag

Human IgG1 Fc tag at the C-terminus

# **Protein Format**

Soluble

#### **Form**

LYOPH

### Reconstitution

Please see Certificate of Analysis for specific instructions.

#### **Buffer**

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4. Normally trehalose is added as protectant before lyophilization.

#### Storage

The product MUST be stored at -70°C or lower upon receipt; -70°C for 3 months under sterile conditions

#### **Target**

#### **Target Protein**

Mucin-1 / MUC-1 (24-380)

#### **Full Name**

mucin 1, cell surface associated

#### Introduction

This gene encodes a membrane-bound protein that is a member of the mucin family. Mucins are O-glycosylated proteins that play an essential role in forming protective mucous barriers on epithelial surfaces. These proteins also play a role in intracellular signaling. This protein is expressed on the apical surface of epithelial cells that line the mucosal surfaces of many different tissues including lung, breast stomach and pancreas. This protein is proteolytically cleaved into alpha and beta subunits that form a heterodimeric complex. The N-terminal alpha subunit functions in cell-adhesion and the C-terminal beta subunit is involved in cell signaling. Overexpression, aberrant intracellular localization, and changes in glycosylation of this protein have been associated with carcinomas. This gene is known to contain a highly polymorphic variable number tandem repeats (VNTR) domain. Alternate splicing results in multiple transcript variants.

#### **Alternative Names**

EMA; MCD; PEM; PUM; KL-6; MAM6; MCKD; PEMT; CD227; H23AG; MCKD1; MUC-1; ADMCKD; ADMCKD1; CA 15-3; MUC-1/X; MUC-1/SEC; mucin-1; H23 antigen; breast carcinoma-associated antigen DF3; cancer antigen 15-3; carcinoma-associated mucin; episialin; krebs von den Lungen-6; mucin 1, transmembrane; peanut-reactive urinary mucin; polymorphic epithelial mucin; tumor associated epithelial mucin; tumor-associated epithelial membrane antigen

Gene ID

4582

**UniProt ID** 

P15941