

## Product Information

### MemDX™ Antibody Discovery - Human ROR1 (30-403) Membrane Protein, Partial, -His tag, [FITC]

Cat. No.: **MP0815F**

This product is for research use only and is not intended for diagnostic use.

This membrane protein is Human ROR1 (30-403). It has been tested in SDS-PAGE, ELISA. We provide this protein to facilitate your membrane protein antibody discovery and development.

#### Product Specifications

##### Host Species

Human

##### Target Protein

ROR1

##### Protein Length

ECD

##### Molecular Weight

The protein has a calculated MW of 43.9 kDa. The protein migrates as 55-66 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

##### Sequence

AA Gln 30 - Glu 403 (Accession # Q01973-1)

#### Product Description

##### Activity

Yes

##### Application

SDS-PAGE, ELISA

##### Expression Systems

HEK293

##### Tag

His Tag at the C-terminus

##### Protein Format

Soluble

##### Form

LYOPH

### **Reconstitution**

Please see Certificate of Analysis for specific instructions.

### **Endotoxin**

<1.0 EU/µg by the LAL method

### **Conjugation**

FITC

### **Purity**

>95% as determined by SDS-PAGE.

### **Buffer**

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4. Normally Trehalose is added as protectant before lyophilization.

### **Storage**

Please protect from light and avoid repeated freeze-thaw cycles.

The product must be protected from light;

2-8 ° C for 12 months in liquid state.

## **Target**

### **Target Protein**

ROR1

### **Full Name**

receptor tyrosine kinase like orphan receptor 1

### **Introduction**

This gene encodes a receptor tyrosine kinase-like orphan receptor that modulates neurite growth in the central nervous system. The encoded protein is a glycosylated type I membrane protein that belongs to the ROR subfamily of cell surface receptors. It is a pseudokinase that lacks catalytic activity and may interact with the non-canonical Wnt signalling pathway. This gene is highly expressed during early embryonic development but expressed at very low levels in adult tissues. Increased expression of this gene is associated with B-cell chronic lymphocytic leukaemia. Alternative splicing results in multiple transcript variants encoding different isoforms.

### **Alternative Names**

NTRKR1; dj537F10.1; inactive tyrosine-protein kinase transmembrane receptor ROR1; neurotrophic tyrosine kinase, receptor-related 1

### **Gene ID**

[4919](#)

### **UniProt ID**

[Q01973](#)