

# Product Information

## MemDX™ Antibody Discovery - Human ROR1 (165-305, Frizzled domain) (165-305)

### Membrane Protein, Partial, -His tag

Cat. No.: **MP0822F**

This product is for research use only and is not intended for diagnostic use.

This membrane protein is Human ROR1 (165-305, Frizzled domain) (308-395). It has been tested in SDS-PAGE. We provide this protein to facilitate your membrane protein antibody discovery and development.

### Product Specifications

#### Host Species

Human

#### Target Protein

ROR1 (165-305, Frizzled domain)

#### Protein Length

ECD

#### Molecular Weight

The protein has a calculated MW of 16.7 kDa. The protein migrates as 19-20 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

#### Sequence

AA Glu 165 - Asp 305 (Accession # Q01973-1)

### Product Description

#### Application

SDS-PAGE

#### Expression Systems

HEK293

#### Tag

His Tag at the C-terminus

#### Protein Format

Soluble

#### Form

LYOPH

#### Reconstitution

Please see Certificate of Analysis for specific instructions.

#### **Endotoxin**

<1.0 EU/µg by the LAL method

#### **Purity**

>95% as determined by SDS-PAGE.

#### **Buffer**

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4. Normally trehalose is added as protectant before lyophilization.

#### **Storage**

Please protect from light and avoid repeated freeze-thaw cycles.

The product must be protected from light;

2-8 ° C for 12 months in liquid state.

### **Target**

#### **Target Protein**

ROR1 (165-305, Frizzled domain)

#### **Full Name**

receptor tyrosine kinase like orphan receptor 1

#### **Introduction**

This gene encodes a receptor tyrosine kinase-like orphan receptor that modulates neurite growth in the central nervous system. The encoded protein is a glycosylated type I membrane protein that belongs to the ROR subfamily of cell surface receptors. It is a pseudokinase that lacks catalytic activity and may interact with the non-canonical Wnt signalling pathway. This gene is highly expressed during early embryonic development but expressed at very low levels in adult tissues. Increased expression of this gene is associated with B-cell chronic lymphocytic leukaemia. Alternative splicing results in multiple transcript variants encoding different isoforms.

#### **Alternative Names**

NTRKR1; dJ537F10.1; inactive tyrosine-protein kinase transmembrane receptor ROR1; neurotrophic tyrosine kinase, receptor-related 1

#### **Gene ID**

[4919](#)

#### **UniProt ID**

[Q01973](#)