

# Product Information

## **MemDX™ Antibody Discovery - Human CD38 (43-300) Membrane Protein, Partial, -hIgG1 Fc tag, [FITC]**

Cat. No.: **MP1114F**

This product is for research use only and is not intended for diagnostic use.

This membrane protein is Human CD38 (43-300). It has been tested in SDS-PAGE, FACS. We provide this protein to facilitate your membrane protein antibody discovery and development.

### Product Specifications

#### Host Species

Human

#### Target Protein

CD38

#### Protein Length

ECD

#### Molecular Weight

The protein has a calculated MW of 56.3 kDa.

#### Sequence

AA Val 43 - Ile 300 (Accession # P28907-1).

### Product Description

#### Activity

Yes

#### Application

SDS-PAGE, FACS

#### Expression Systems

HEK293

#### Tag

Human IgG1 Fc tag at the C-terminus

#### Protein Format

Soluble

#### Form

LYOPH

### Reconstitution

Please see Certificate of Analysis for specific instructions.

### Conjugation

FITC

### Buffer

Please contact us for detailed information.

Contact us for customized product form or formulation.

### Storage

Stored at lyophilized form at -20°C or lower. Avoid repeated freeze-thaw cycles.

The antigen can be stable for 12 months in lyophilized form after storage at -20°C to -80°C, 3 months under sterile conditions after reconstitution after storage at -80°C.

## Target

### Target Protein

CD38

### Full Name

CD38 molecule

### Introduction

The protein encoded by this gene is a non-lineage-restricted, type II transmembrane glycoprotein that synthesizes and hydrolyzes cyclic adenosine 5'-diphosphate-ribose, an intracellular calcium ion mobilizing messenger. The release of soluble protein and the ability of membrane-bound protein to become internalized indicate both extracellular and intracellular functions for the protein. This protein has an N-terminal cytoplasmic tail, a single membrane-spanning domain, and a C-terminal extracellular region with four N-glycosylation sites. Crystal structure analysis demonstrates that the functional molecule is a dimer, with the central portion containing the catalytic site. It is used as a prognostic marker for patients with chronic lymphocytic leukemia. Alternative splicing results in multiple transcript variants.

### Alternative Names

ADPRC1; ADPRC 1; ADP-ribosyl cyclase/cyclic ADP-ribose hydrolase 1; 2'-phospho-ADP-ribosyl cyclase; 2'-phospho-cyclic-ADP-ribose transferase; ADP-ribosyl cyclase 1; CD38 antigen (p45); NAD(+) nucleosidase; cluster of differentiation 38; cyclic ADP-ribose hydrolase 1; ecto-nicotinamide adenine dinucleotide glycohydrolase

### Gene ID

[952](#)

### UniProt ID

[P28907](#)