

## Product Information

### **MemDX™ Antibody Discovery - Human CD40 / TNFRSF (21-193) Membrane Protein, Partial, -hIgG1 Fc -Avi tag, [Biotin]**

Cat. No.: **MP1132F**

This product is for research use only and is not intended for diagnostic use.

This membrane protein is Human CD40 / TNFRSF (21-193). It has been tested in SDS-PAGE, ELISA, SEC-SEC-MALS, FACS. We provide this protein to facilitate your membrane protein antibody discovery and development.

#### Product Specifications

##### Host Species

Human

##### Target Protein

CD40 / TNFRSF

##### Protein Length

ECD

##### Molecular Weight

The protein has a calculated MW of 48.1 kDa. The protein migrates as 60-66 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

##### Sequence

AA Glu 21 - Arg 193 (Accession # P25942-1).

#### Product Description

##### Activity

Yes

##### Application

SDS-PAGE, ELISA, SEC-SEC-MALS, FACS

##### Expression Systems

HEK293

##### Tag

Human IgG1 Fc tag at the C-terminus, followed by a Avi tag

##### Protein Format

Soluble

##### Form

LYOPH

### Reconstitution

Please see Certificate of Analysis for specific instructions.

### Endotoxin

<1.0 EU/μg by the LAL method

### Conjugation

Biotin

### Purity

>95% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

### Buffer

Lyophilized from 0.22 μm filtered solution in Tris with Glycine, Arginine and NaCl, pH7.5. Normally trehalose is added as protectant before lyophilization.

### Storage

Stored at lyophilized form at -20°C or lower. Avoid repeated freeze-thaw cycles.

The antigen can be stable for 12 months in lyophilized form after storage at -20°C to -80°C, 3 months under sterile conditions after reconstitution after storage at -80°C.

## Target

### Target Protein

CD40 / TNFRSF

### Full Name

CD40 molecule

### Introduction

This gene is a member of the TNF-receptor superfamily. The encoded protein is a receptor on antigen-presenting cells of the immune system and is essential for mediating a broad variety of immune and inflammatory responses including T cell-dependent immunoglobulin class switching, memory B cell development, and germinal center formation. AT-hook transcription factor AKNA is reported to coordinately regulate the expression of this receptor and its ligand, which may be important for homotypic cell interactions. Adaptor protein TNFR2 interacts with this receptor and serves as a mediator of the signal transduction. The interaction of this receptor and its ligand is found to be necessary for amyloid-beta-induced microglial activation, and thus is thought to be an early event in Alzheimer disease pathogenesis. Mutations affecting this gene are the cause of autosomal recessive hyper-IgM immunodeficiency type 3 (HIGM3). Multiple alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported.

### Alternative Names

CD40, CD40 molecule, TNF receptor superfamily member 5, TNFRSF5, tumor necrosis factor receptor superfamily, member 5, tumor necrosis factor receptor superfamily member 5, Bp50, p50, CD40L receptor, CD40 type II isoform, B cell-associated molecule, B cell surface antigen CD40, B-cell surface antigen CD40, CD40 antigen (TNF receptor superfamily member 5), tumor necrosis factor receptor superfamily, member 5, nerve growth factor receptor-related B-lymphocyte activation molecule, CDW40, TNFRSF5, MGC9013,

### Gene ID

[958](#)

### UniProt ID

[P25942](#)