

Product Information

MemDX™ Antibody Discovery - Human CD4 (26-396) Membrane Protein, Partial, -hIgG1 Fc tag, [FITC]

Cat. No.: **MP1260F**

This product is for research use only and is not intended for diagnostic use.

This membrane protein is Human CD4 (26-396). It has been tested in SDS-PAGE, ELISA, FACS. We provide this protein to facilitate your membrane protein antibody discovery and development.

Product Specifications

Host Species

Human

Target Protein

CD4

Protein Length

ECD

Molecular Weight

The protein has a calculated MW of 67.7 kDa. The protein migrates as 67-80 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Sequence

AA Lys 26 - Pro 396 (Accession # AAH25782). .

Product Description

Activity

Yes

Application

SDS-PAGE, ELISA, FACS

Expression Systems

HEK293

Tag

Human IgG1 Fc tag at the C-terminus

Protein Format

Soluble

Form

LYOPH

Reconstitution

Please see Certificate of Analysis for specific instructions.

Endotoxin

<1.0 EU/μg by the LAL method

Conjugation

FITC

Purity

>95% as determined by SDS-PAGE.

Buffer

Lyophilized from 0.22 μm filtered solution in PBS, pH7.4. Normally trehalose is added as protectant before lyophilization.

Storage

Stored at lyophilized form at -20°C or lower. Avoid repeated freeze-thaw cycles.

The antigen can be stable for 12 months in lyophilized form after storage at -20°C to -80°C, 3 months under sterile conditions after reconstitution after storage at -80°C.

Target

Target Protein

CD4

Full Name

CD4 molecule

Introduction

This gene encodes the CD4 membrane glycoprotein of T lymphocytes. The CD4 antigen acts as a coreceptor with the T-cell receptor on the T lymphocyte to recognize antigens displayed by an antigen presenting cell in the context of class II MHC molecules. The CD4 antigen is also a primary receptor for entry of the human immunodeficiency virus through interactions with the HIV Env gp120 subunit. This gene is expressed not only in T lymphocytes, but also in B cells, macrophages, granulocytes, as well as in various regions of the brain. The protein functions to initiate or augment the early phase of T-cell activation, and may function as an important mediator of indirect neuronal damage in infectious and immune-mediated diseases of the central nervous system. Multiple alternatively spliced transcript variants encoding different isoforms have been identified in this gene.

Alternative Names

CD4mut; T-cell surface glycoprotein CD4; CD4 antigen (p55); CD4 receptor; T-cell surface antigen T4/Leu-3

Gene ID

[920](#)

UniProt ID

[P01730](#)