

Product Information

MemDX™ Membrane Protein Human CCR1 (C-C motif chemokine receptor 1) for Antibody

Discovery

Cat. No.: **MP1329J**

This product is for research use only and is not intended for diagnostic use.

This product is a 41.2 kDa Human CCR1 membrane protein expressed in E.coli. The protein is for research use only and is not approved for use in humans or in clinical diagnosis.

Product Specifications

Host Species

Human

Target Protein

CCR1

Protein Length

Full-length

Protein Class

GPCR

Molecular Weight

41.2 kDa

TMD

7

Sequence

METPNTTEDYDTTTEFDYGDATPCQKVNERAFGAQLLPPLYSLVFVIGLVGNILVVLVLV
QYKRLKNMTSIYLLNLAISDLLFLFTLPFWIDYKLKDDWVFGDAMCKILSGFYTTGLYSE
IFFIILLTIDRYLAIVHAVFALRARTVTFGVITSIIWALAILASMPGLYFSKTQWEFTH
HTCSLHFPHESLREWKLFGALKNLFGVLPLLVMIICYTGIIKILLRRPNEKKSKAVRL
IFVIMIIFFLFWTPYNLTILISVFQDFLFTHCEQSRHLDLAVQVTEVIAYTHCCVNPVI
YAFVGERFRKYLRQLFHRRVAVHLVKWLPFLSVDRLERVSSTSPSTGEHELSAGF

Product Description

Expression Systems

E.coli

Tag

N-His or Tag-Free

Form

Lyophilized powder

Reconstitution

Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration).

Purity

>85% as determined by SDS-PAGE

Buffer

Tris/PBS-based buffer, 6% Trehalose, pH 8.0

Storage

Store at +4°C for up to one week or several months at -80°C

Target**Target Protein**

CCR1

Full Name

C-C motif chemokine receptor 1

Introduction

This gene encodes a member of the beta chemokine receptor family, which is predicted to be a seven transmembrane protein similar to G protein-coupled receptors. The ligands of this receptor include macrophage inflammatory protein 1 alpha (MIP-1 alpha), regulated on activation normal T expressed and secreted protein (RANTES), monocyte chemoattractant protein 3 (MCP-3), and myeloid progenitor inhibitory factor-1 (MPIF-1). Chemokines and their receptors mediated signal transduction are critical for the recruitment of effector immune cells to the site of inflammation. Knockout studies of the mouse homolog suggested the roles of this gene in host protection from inflammatory response, and susceptibility to virus and parasite. This gene and other chemokine receptor genes, including CCR2, CCRL2, CCR3, CCR5 and CXCR1, are found to form a gene cluster on chromosome 3p.

Alternative Names

C C chemokine receptor type 1; C C CKR 1; C-C chemokine receptor type 1; C-C CKR-1; CC CKR 1; CC-CKR-1; CCR 1; CCR-1; CCR1; CCR1_HUMAN; CD 191; CD191; CD191 antigen; Chemokine (C C motif) receptor 1; Chemokine C C motif receptor 1; CKR 1; CKR1; CMKBR 1; CMKBR1; CMKR 1; CMKR1; HM145; LD78 receptor; Macrophage inflammatory protein 1 alpha /Rantes receptor; Macrophage inflammatory protein 1 alpha receptor; Macrophage inflammatory protein 1-alpha receptor; Mip 1a R; MIP 1alpha R; MIP-1alpha-R; MIP1aR; RANTES R; RANTES receptor; RANTES-R; SCYAR1

Gene ID

[1230](#)

UniProt ID

[P32246](#)