

Product Information

MemDX™ Antibody Discovery - Cynomolgus/Rhesus macaque FcRn (FCGRT & B2M) (24-297(FCGRT)&21-119(B2M)) Membrane Protein, Partial

Cat. No.: **MP1434F**

This product is for research use only and is not intended for diagnostic use.

This membrane protein is Cynomolgus/Rhesus macaque FcRn (FCGRT & B2M) (24-297(FCGRT)&21-119(B2M)). It has been tested in SDS-PAGE, SPR, BLI. We provide this protein to facilitate your membrane protein antibody discovery and development.

Product Specifications

Host Species

Cynomolgus / Rhesus macaque

Target Protein

FcRn (FCGRT & B2M)

Protein Length

ECD

Molecular Weight

32.3 kDa (FCGRT) and 13.1 kDa (B2M)

Sequence

AA Ala 24 - Ser 297 (FCGRT) & Ile 21 - Met 119 (B2M) (Accession # Q8SPV9-1 (FCGRT) & Q8SPW0-1 (B2M)).

Product Description

Activity

Yes

Application

SDS-PAGE, SPR, BLI

Expression Systems

HEK293

Protein Format

Soluble

Form

LYOPH

Reconstitution

Please see Certificate of Analysis for specific instructions.

Endotoxin

<1.0 EU/μg by the LAL method

Purity

>95% as determined by SDS-PAGE

Buffer

Lyophilized from 0.22 μm filtered solution in PBS, pH7.4. Normally trehalose is added as protectant before lyophilization.

Storage

Stored at lyophilized form at -20°C or lower. Avoid repeated freeze-thaw cycles.

The antigen can be stable for 12 months in lyophilized form after storage at -20°C to -80°C, 3 months under sterile conditions after reconstitution after storage at -80°C.

Target**Target Protein**

FcRn (FCGRT & B2M)

Full Name

Fc fragment of IgG receptor and transporter & beta-2-microglobulin

Introduction

This gene encodes a receptor that binds the Fc region of monomeric immunoglobulin G. The encoded protein transfers immunoglobulin G antibodies from mother to fetus across the placenta. This protein also binds immunoglobulin G to protect the antibody from degradation. Alternative splicing results in multiple transcript variants.

Alternative Names

FCRN, alpha-chain, IgG receptor FcRn large subunit p51, Fc fragment of IgG, receptor, transporter, alpha, FcRn alpha chain, IgG Fc fragment receptor transporter alpha chain, heavy chain of the major histocompatibility complex class I-like Fc receptor, immunoglobulin receptor, intestinal, heavy chain, major histocompatibility complex class I-like Fc receptor, neonatal Fc receptor, neonatal Fc-receptor for Ig, transmembrane alpha chain of the neonatal receptor

Gene ID

[102128913](#); [712428](#)

UniProt ID

[Q8SPV9](#); [Q6V7J5](#)