SUITE 203, 17 Ramsey Road, Shirley, NY 11967, USA
Tel: 1-631-416-1478 Fax: 1-631-207-8356
Email: info@creative-biolabs.com
www.creative-biolabs.com

Product Information

MemDX™ Membrane Protein Human UCP3 (Uncoupling protein 3) Full Length

Cat. No.: **MPC2644K**
This product is for research use only and is not intended for diagnostic use.

This product is a made-to-order Human UCP3 membrane protein expressed in HEK293. The protein is for research use only and is not approved for use in humans or in clinical diagnosis.

Product Specifications

Host Species
Human

Target Protein
UCP3

Protein Length
Full length

Protein Class
Transporter

TMD
6

Sequence
MVGLKPSDVPPTMAVKFLGAGTAACFADLVTFPLDTAKVRLQIQGENQAV
QTARLVQYRGVLGTILTMVRTEGPCSPYNGLVAGLQRQMSFASIRIGLYD
SVKQVYTPKGADNSSLTTRILAGCTTGAMAVTCAQPTDVVKVRFQASIHL
GPSRSDRKYSGTMDAYRTIAREEGVRGLWKGTLPNIMRNAIVNCAEVVTY
DILKEKLLDYHLLTDNFPCHFVSAFGAGFCATVVASPVDVVKTRYMNSPP
GQYFSPLDCMIKMVAQEGPTAFYKGFTPSFLRLGSWNVVMFVTYEQLKRA
LMKVQMLRESPF

Product Description

Expression Systems
HEK293

Tag
Based on specific requirements

Protein Format
Detergent or based on specific requirements (Detergent, Liposome, Nanodisc, Polymer, VLP)

Form

Liquid

Storage

Aliquot and store at -20°C or lower. For long term storage, we recommend to store at -72°C or lower. Avoid freeze/thaw cycles.

Target

Target Protein

UCP3

Full Name

Uncoupling protein 3

Introduction

Mitochondrial uncoupling proteins (UCP) are members of the larger family of mitochondrial anion carrier proteins (MACP). UCPs separate oxidative phosphorylation from ATP synthesis with energy dissipated as heat, also referred to as the mitochondrial proton leak. UCPs facilitate the transfer of anions from the inner to the outer mitochondrial membrane and the return transfer of protons from the outer to the inner mitochondrial membrane. They also reduce the mitochondrial membrane potential in mammalian cells. The different UCPs have tissue-specific expression; this gene is primarily expressed in skeletal muscle. This gene's protein product is postulated to protect mitochondria against lipid-induced oxidative stress. Expression levels of this gene increase when fatty acid supplies to mitochondria exceed their oxidation capacity and the protein enables the export of fatty acids from mitochondria. UCPs contain the three solcar protein domains typically found in MACPs. Two splice variants have been found for this gene.

Alternative Names

UCP3; SLC25A9; mitochondrial uncoupling protein 3; solute carrier family 25 member 9; uncoupling protein 3 (mitochondrial, proton carrier); Uncoupling protein 3

Gene ID

7352

UniProt ID

P55916

SUITE 203, 17 Ramsey Road, Shirley, NY 11967, USA
Tel: 1-631-416-1478 Fax: 1-631-207-8356
Email: info@creative-biolabs.com