

Product Information

MemDX™ Membrane Protein Human OR10AC1 (Olfactory receptor family 10 subfamily AC member 1 (gene/pseudogene)) Full Length

Cat. No.: **MPC2705K**

This product is for research use only and is not intended for diagnostic use.

This product is a made-to-order Human OR10AC1 membrane protein expressed in Baculovirus/Insect expression system. The protein is for research use only and is not approved for use in humans or in clinical diagnosis.

Product Specifications

Host Species

Human

Target Protein

OR10AC1

Protein Length

Full length

Protein Class

GPCR

TMD

7

Sequence

MDSPSNATVPCGFLLQGFSEFPHLRPVFLLLLGVHLATLGGNLLILVAV
ASMPSRQPMLLFLCQLSAIELCYTLVVVPRSLVDLSTPGHRRGSPISFLS
CAFQMFMFVALGGAECFLAAMAYDRYVAICHPLRYAAVVTPGLCARLAL
ACCLRGLAVSVGLTVAIFHLPFCGSRLLLHFFCDITALLHLACTRSYADE
LPLLGACLVLLLLPSVLILASYGAIAAALRRLRCPKGRGKAASTCALHLA
VTFLHYGCATFMYVRPRASYSRDLRTLALVYTNVTPLLCLPIYSLRNRE
ITAALSRVLGRRRPGQAPGGDLREL

Product Description

Expression Systems

Baculovirus/Insect expression system

Tag

Based on specific requirements

Protein Format

Detergent or based on specific requirements (Detergent, Liposome, Nanodisc, Polymer, VLP)

Form

Liquid

Storage

Aliquot and store at -20°C or lower. For long term storage, we recommend to store at -72°C or lower. Avoid freeze/thaw cycles.

Target**Target Protein**

OR10AC1

Full Name

Olfactory receptor family 10 subfamily AC member 1 (gene/pseudogene)

Introduction

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. This olfactory receptor gene is a segregating pseudogene, where some individuals have an allele that encodes a functional olfactory receptor, while other individuals have an allele encoding a protein that is predicted to be non-functional.

Alternative Names

OR10AC1; OR10AC1P; olfactory receptor 10AC1; Olfactory receptor OR7-5; Putative olfactory receptor 10AC1; olfactory receptor, family 10, subfamily AC, member 1 pseudogene; olfactory receptor, family 10, subfamily W, member 1 pseudogene; seven transmembrane helix receptor; Olfactory receptor family 10 subfamily AC member 1 (gene/pseudogene)

Gene ID

[392133](#)

UniProt ID

[Q8NH08](#)