

Product Information

MemDX™ Human ARV7 HEK293 Cell Line

Cat. No.: **S01YF-0123-KX105**

This product is for research use only and is not intended for diagnostic use.

Product Information

Target Protein

ARV7

Target Protein Species

Human

Accession Number

NM_001348061.1

Protein Tag

Tag-free

Host Cell Type

HEK293

Target Classification

Nuclear Receptor

Target Family

Nuclear Receptor

Target Research Area

Inflammation Research; Reproductive Research

Related Diseases

Androgen Insensitivity; Partial and Spinal And Bulbar Muscular Atrophy

Product Properties

Morphology

Fibroblastoid cells growing as a monolayer

Assay Types

Drug screening and biological assays

Resistance

Puromycin

Mycoplasma Testing

Negative

Sterility Testing

10 passages

Biosafety Level

Level 1

Activity

Yes

Quantity

5x10⁶ cells

Form

Frozen cells

Freeze Medium

70% DMEM + 20% FBS + 10% DMSO

Culture Medium

DMEM + 10% FBS + 1µg/mL Puromycin

Selective Antibiotic(s)

Regular antibiotics active against mycoplasmas, bacteria and fungi.

Handling Notes

Frozen cells should be thawed immediately upon receipt and grown according to handling procedure to ensure cell viability and proper assay performance.

Note: Do not freeze the cells upon receipt as it may result in irreversible damage to the cell line.

Disclaimer: We cannot guarantee cell viability if the cells are not thawed immediately upon receipt and grown according to handling procedure.

Incubation

37°C with 5% CO₂

Applications

Drug screening and biological assays

Application Notes

Cells were plated in a 384-well plate and incubated overnight at 37°C and 5% CO₂ to allow the cells to attach and grow. Cells were then stimulated with a control for high-throughput drugs screening and functional assays.

Use Restrictions

These cells are distributed for research use only.

Shipping

Dry ice

Storage

Liquid nitrogen

Target**Full Name**

Introduction

The androgen receptor gene is more than 90 kb long and codes for a protein that has 3 major functional domains: the N-terminal domain, DNA-binding domain, and androgen-binding domain. The protein functions as a steroid-hormone activated transcription factor. Upon binding the hormone ligand, the receptor dissociates from accessory proteins, translocates into the nucleus, dimerizes, and then stimulates transcription of androgen responsive genes. This gene contains 2 polymorphic trinucleotide repeat segments that encode polyglutamine and polyglycine tracts in the N-terminal transactivation domain of its protein. Expansion of the polyglutamine tract from the normal 9-34 repeats to the pathogenic 38-62 repeats causes spinal bulbar muscular atrophy (SBMA, also known as Kennedy's disease). Mutations in this gene are also associated with complete androgen insensitivity (CAIS). Alternative splicing results in multiple transcript variants encoding different isoforms.

Alternative Names

KD; AIS; AR8; TFM; DHTR; SBMA; HYSP1; NR3C4; SMAX1; HUMARA; androgen receptor; dihydrotestosterone receptor; nuclear receptor subfamily 3 group C member 4

Gene ID

[367](#)

UniProt ID

[P10275](#)