

# Product Information

## MemDX™ Human LILRB2 CT26 Cell Line

Cat. No.: **S01YF-0123-KX181**

This product is for research use only and is not intended for diagnostic use.

### Product Information

#### Target Protein

LILRB2

#### Target Protein Species

Human

#### Accession Number

NM\_005874.5

#### Protein Tag

Tag-free

#### Host Cell Type

CT26

#### Target Classification

Oncology

#### Target Family

Oncology

#### Target Research Area

CNS Research; Immunology Research; Infectious Research

#### Related Diseases

Hymenolepiasis

### Product Properties

#### Morphology

Epithelial

#### Assay Types

Drug screening and biological assays

#### Resistance

Puromycin

#### Mycoplasma Testing

Negative

**Sterility Testing**

10 passages

**Biosafety Level**

Level 1

**Activity**

Yes

**Quantity**

5x10<sup>6</sup> cells

**Form**

Frozen cells

**Freeze Medium**

70% DMEM + 20% FBS + 10% DMSO

**Culture Medium**

DMEM + 10% FBS + 10µg/mL Puromycin

**Selective Antibiotic(s)**

Regular antibiotics active against mycoplasmas, bacteria and fungi.

**Handling Notes**

Frozen cells should be thawed immediately upon receipt and grown according to handling procedure to ensure cell viability and proper assay performance.

Note: Do not freeze the cells upon receipt as it may result in irreversible damage to the cell line.

Disclaimer: We cannot guarantee cell viability if the cells are not thawed immediately upon receipt and grown according to handling procedure.

**Incubation**

37°C with 5% CO<sub>2</sub>

**Applications**

Drug screening and biological assays

**Application Notes**

Cells were plated in a 384-well plate and incubated overnight at 37°C and 5% CO<sub>2</sub> to allow the cells to attach and grow. Cells were then stimulated with a control for high-throughput drugs screening and functional assays.

**Use Restrictions**

These cells are distributed for research use only.

**Shipping**

Dry ice

**Storage**

Liquid nitrogen

**Target****Full Name**

### Introduction

This gene is a member of the leukocyte immunoglobulin-like receptor (LIR) family, which is found in a gene cluster at chromosomal region 19q13.4. The encoded protein belongs to the subfamily B class of LIR receptors which contain two or four extracellular immunoglobulin domains, a transmembrane domain, and two to four cytoplasmic immunoreceptor tyrosine-based inhibitory motifs (ITIMs). The receptor is expressed on immune cells where it binds to MHC class I molecules on antigen-presenting cells and transduces a negative signal that inhibits stimulation of an immune response. It is thought to control inflammatory responses and cytotoxicity to help focus the immune response and limit autoreactivity. Multiple transcript variants encoding different isoforms have been found for this gene.

### Alternative Names

LILRB2; ILT4; LIR2; CD85D; ILT-4; LIR-2; MIR10; MIR-10; leukocyte immunoglobulin-like receptor subfamily B member 2; CD85 antigen-like family member D; Ig-like transcript 4; leucocyte Ig-like receptor B2; leukocyte immunoglobulin-like receptor, subfamily B (with TM and ITIM domains), member 2; monocyte/macrophage immunoglobulin-like receptor 10; myeloid inhibitory receptor 10; Leukocyte immunoglobulin like receptor B2

### Gene ID

[10288](#)

### UniProt ID

[Q8N423](#)