

# Product Information

## MemDX™ Human TNFRSF1B MC-38 Cell Line

Cat. No.: **S01YF-0324-KX134**

This product is for research use only and is not intended for diagnostic use.

### Product Information

#### Target Protein

TNFRSF1B

#### Target Protein Species

Human

#### Host Cell Type

MC-38

#### Target Classification

Others

#### Target Family

Others

#### Target Research Area

Infectious Research

#### Related Diseases

Mycosis Fungoides; Sezary's Disease

### Product Properties

#### Assay Types

Functional assay and biological assay

#### Mycoplasma Testing

Negative

#### Biosafety Level

Level 1

#### Activity

Yes

#### Form

Frozen cells

#### Freeze Medium

70% RPMI 1640 + 20% FBS + 10% DMSO

### Culture Medium

RPMI 1640 + 10% FBS

### Selective Antibiotic(s)

Regular antibiotics active against mycoplasmas, bacteria and fungi.

### Handling Notes

Frozen cells should be thawed immediately upon receipt and grown according to handling procedure to ensure cell viability and proper assay performance.

Note: Do not freeze the cells upon receipt as it may result in irreversible damage to the cell line.

Disclaimer: We cannot guarantee cell viability if the cells are not thawed immediately upon receipt and grown according to handling procedure.

### Incubation

37°C with 5% CO<sub>2</sub>

### Applications

Drug screening and biological assays

### Application Notes

Cells were plated in a 384-well plate and incubated overnight at 37°C and 5% CO<sub>2</sub> to allow the cells to attach and grow. Cells were then stimulated with a control for high-throughput drugs screening and functional assays.

### Use Restrictions

These cells are distributed for research use only.

### Shipping

Dry ice

### Storage

Liquid nitrogen

## Target

### Full Name

TNF receptor superfamily member 1B

### Introduction

The protein encoded by this gene is a member of the TNF-receptor superfamily. This protein and TNF-receptor 1 form a heterocomplex that mediates the recruitment of two anti-apoptotic proteins, c-IAP1 and c-IAP2, which possess E3 ubiquitin ligase activity. The function of IAPs in TNF-receptor signalling is unknown, however, c-IAP1 is thought to potentiate TNF-induced apoptosis by the ubiquitination and degradation of TNF-receptor-associated factor 2, which mediates anti-apoptotic signals. Knockout studies in mice also suggest a role of this protein in protecting neurons from apoptosis by stimulating antioxidative pathways.

### Alternative Names

TNFRSF1B; p75; TBPII; TNFBR; TNFR2; CD120b; TNFR1B; TNFR80; TNF-R75; p75TNFR; TNF-R-II; tumor necrosis factor receptor superfamily member 1B; TNF-R2; TNF-RII; p75 TNF receptor; p80 TNF-alpha receptor; soluble TNFR1B variant 1; tumor necrosis factor beta receptor; tumor necrosis factor binding protein 2; tumor necrosis factor receptor 2; tumor necrosis factor receptor type II; TNF receptor superfamily member 1B

### Gene ID

7133

**UniProt ID**

P20333