SUITE 203, 17 Ramsey Road, Shirley, NY 11967, USA
Tel: 1-631-416-1478 Fax: 1-631-207-8356
Email: info@creative-biolabs.com
www.creative-biolabs.com

Product Information

MemDX™ Human TNFRSF1B CT26 Cell Line

Cat. No.: **S01YF-0324-KX66**
This product is for research use only and is not intended for diagnostic use.

Product Information

Target Protein
TNFRSF1B

Target Protein Species
Human

Host Cell Type
CT26

Target Classification
Others

Target Family
Others

Target Research Area
Infectious Research

Related Diseases
Mycosis Fungoides; Sezary's Disease

Product Properties

Assay Types
Functional assay and biological assay

Mycoplasma Testing
Negative

Biosafety Level
Level 1

Activity
Yes

Form
Frozen cells

Freeze Medium

DMEM + 20% FBS + 10% DMSO

Culture Medium

DMEM + 10% FBS

Selective Antibiotic(s)

Regular antibiotics active against mycoplasmas, bacteria and fungi.

Handling Notes

Frozen cells should be thawed immediately upon receipt and grown according to handling procedure to ensure cell viability and proper assay performance.
Note: Do not freeze the cells upon receipt as it may result in irreversible damage to the cell line.
Disclaimer: We cannot guarantee cell viability if the cells are not thawed immediately upon receipt and grown according to handling procedure.

Incubation

37°C with 5% CO_2

Applications

Drug screening and biological assays

Application Notes

Cells were plated in a 384-well plate and incubated overnight at 37°C and 5% CO_2 to allow the cells to attach and grow. Cells were then stimulated with a control for high-throughput drugs screening andfunctional assays.

Use Restrictions

These cells are distributed for research use only.

Shipping

Dry ice

Storage

Liquid nitrogen

Target

Full Name

TNF receptor superfamily member 1B

Introduction

The protein encoded by this gene is a member of the TNF-receptor superfamily. This protein and TNF-receptor 1 form a heterocomplex that mediates the recruitment of two anti-apoptotic proteins, c-IAP1 and c-IAP2, which possess E3 ubiquitin ligase activity. The function of IAPs in TNF-receptor signalling is unknown, however, c-IAP1 is thought to potentiate TNF-induced apoptosis by the ubiquitination and degradation of TNF-receptor-associated factor 2, which mediates anti-apoptotic signals. Knockout studies in mice also suggest a role of this protein in protecting neurons from apoptosis by stimulating antioxidative pathways.

Alternative Names

TNFRSF1B; p75; TBPII; TNFBR; TNFR2; CD120b; TNFR1B; TNFR80; TNF-R75; p75TNFR; TNF-R-II; tumor necrosis factor receptor superfamily member 1B; TNF-R2; TNF-RII; p75 TNF receptor; p80 TNF-alpha receptor; soluble TNFR1B variant 1; tumor necrosis factor beta receptor; tumor necrosis factor binding protein 2; tumor necrosis factor receptor 2; tumor necrosis factor receptor type II; TNF receptor superfamily member 1B

Gene ID

7133

UniProt ID

P20333

SUITE 203, 17 Ramsey Road, Shirley, NY 11967, USA
Tel: 1-631-416-1478 Fax: 1-631-207-8356
Email: info@creative-biolabs.com