

Product Information

MemDX™ Human CD38 EL4 Cell Line

Cat. No.: **S01YF-0324-KX71**

This product is for research use only and is not intended for diagnostic use.

Product Information

Target Protein

CD38

Target Protein Species

Human

Host Cell Type

EL4

Target Classification

Immune Checkpoint

Target Family

Immune Checkpoint

Target Research Area

Autoimmune Research; Immunology Research

Related Diseases

Leukemia; Prolymphocytic Leukemia

Product Properties

Assay Types

Functional assay and biological assay

Mycoplasma Testing

Negative

Biosafety Level

Level 1

Activity

Yes

Form

Frozen cells

Freeze Medium

DMEM + 20% FBS + 10% DMSO

Culture Medium

DMEM + 10% FBS

Selective Antibiotic(s)

Regular antibiotics active against mycoplasmas, bacteria and fungi.

Handling Notes

Frozen cells should be thawed immediately upon receipt and grown according to handling procedure to ensure cell viability and proper assay performance.

Note: Do not freeze the cells upon receipt as it may result in irreversible damage to the cell line.

Disclaimer: We cannot guarantee cell viability if the cells are not thawed immediately upon receipt and grown according to handling procedure.

Incubation

37°C with 5% CO₂

Applications

Drug screening and biological assays

Application Notes

Cells were plated in a 384-well plate and incubated overnight at 37°C and 5% CO₂ to allow the cells to attach and grow. Cells were then stimulated with a control for high-throughput drugs screening and functional assays.

Use Restrictions

These cells are distributed for research use only.

Shipping

Dry ice

Storage

Liquid nitrogen

Target

Full Name

CD38 molecule

Introduction

The protein encoded by this gene is a non-lineage-restricted, type II transmembrane glycoprotein that synthesizes and hydrolyzes cyclic adenosine 5'-diphosphate-ribose, an intracellular calcium ion mobilizing messenger. The release of soluble protein and the ability of membrane-bound protein to become internalized indicate both extracellular and intracellular functions for the protein. This protein has an N-terminal cytoplasmic tail, a single membrane-spanning domain, and a C-terminal extracellular region with four N-glycosylation sites. Crystal structure analysis demonstrates that the functional molecule is a dimer, with the central portion containing the catalytic site. It is used as a prognostic marker for patients with chronic lymphocytic leukemia. Alternative splicing results in multiple transcript variants.

Alternative Names

CD38; ADPRC1; ADPRC 1; ADP-ribosyl cyclase/cyclic ADP-ribose hydrolase 1; 2'-phospho-ADP-ribosyl cyclase; 2'-phospho-cyclic-ADP-ribose transferase; ADP-ribosyl cyclase 1; CD38 antigen (p45); NAD(+) nucleosidase; cluster of differentiation 38; cyclic ADP-ribose hydrolase 1; ecto-nicotinamide adenine dinucleotide glycohydrolase; CD38 molecule

Gene ID

952

UniProt ID

P28907